

60th Annual Scientific Session & Expo

E474

JACC April 5, 2011
Volume 57, Issue 14CONGENITAL CARDIOLOGY SOLUTIONS
(ADULT CONGENITAL AND PEDIATRIC CARDIOLOGY)

EXPERIMENTAL FETAL DUCTAL CONSTRICTION AFTER MATERNAL INGESTION OF GREEN TEA IN LATE PREGNANCY

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Tuesday, April 05, 2011, 9:30 a.m.-10:45 a.m.

Session Title: Pediatric Cardiology: Imaging

Abstract Category: 41 Pediatric Cardiology

Session-Poster Board Number: 1173-438

Authors: *Paulo Zielinsky, Antônio L. Piccoli, Luiz Henrique S. Nicoloso, João L L Manica, Marinez Barra, Marcelo M. Alievi, Izabele Vian, Patrícia E. Pizzato, Luciano P. Bender, Marcelo Pizzato, Fetal Cardiology Unit, Instituto de Cardiologia do Rio Grande do Sul/ FUC (IC/FUC), Porto Alegre, Brazil*

Background: Maternal ingestion of inhibitors of prostaglandin biosynthesis in late pregnancy may cause fetal ductal constriction. Green tea is rich in polyphenols, whose anti-inflammatory effects depend on prostaglandin inhibition. This study was designed to test the hypothesis that experimental maternal intake of green tea in late pregnancy causes fetal ductus arteriosus constriction.

Methods: Twelve fetal lambs (pregnancy >120 days) were assessed before and after maternal administration of green tea to eight and water to four (controls) as the only source of liquid. After 1 week, all specimens exposed to green tea showed at autopsy dilated and hypertrophic right ventricles, not present in control fetuses.

Results: Histological analysis showed a significantly larger mean thickness of the medial avascular zone of the ductus arteriosus in fetuses exposed to green tea than in controls ($747.6 \pm 214.6 \mu\text{m}$ versus $255.3 \pm 97.9 \mu\text{m}$, $p < 0.001$). Fetal Doppler echocardiography also showed evidences of constriction of ductus arteriosus in all fetuses exposed to maternal ingestion of green tea, with increase in mean systolic velocity ($0.70 \pm 0.19 \text{ m/s}$ to $0.92 \pm 0.15 \text{ m/s}$, $P = 0.001$ and mean diastolic velocity ($0.19 \pm 0.05 \text{ m/s}$ to $0.31 \pm 0.01 \text{ m/s}$, $P < 0.001$), decrease of pulsatility index (2.2 ± 0.4 to 1.8 ± 0.3 , $p = 0.003$) and increase of mean RV/LV ratio (0.89 ± 0.14 to 1.43 ± 0.23 , $P < 0.001$). In the four control fetuses receiving only water, there were no significant changes.

Conclusions: This study shows a cause and effect relationship between experimental maternal exposure of green tea and fetal ductus arteriosus constriction in late pregnancy.